

PART SEVENTEEN ANSWERS VOCABULARY PRACTICE

Replace words in italics with the synonym from the table below

Bruxism (4)	Mobility (1)	Extraction (8)	Stabilize (6)
Clenching (7)	gum disease (2)	Stretching (5)	Enlarged (3)

1. Certain amount of tooth *looseness* is always present and it is natural, as the teeth are not fixed tight in their sockets.
2. Though the most common reason for a loose tooth in adults is *periodontal disease*, trauma, bite injury or tumor can also cause teeth mobility.
3. The tissues around a loose tooth become red, *swollen* and damaged.
4. For most adults, oral trauma which leads to loose tooth, is usually caused by *nighttime tooth grinding*.
5. One may even grind one's teeth back and forth, further *lengthening* the ligaments.
6. The first step to treat loose tooth will be to give the tooth some extra support in order *to firm it up*.
7. Sometimes nightguard is used to prevent a patient from fully *tighten* his/her jaws during the night.
8. If the tooth is beyond saving, a tooth *removal* may be necessary.

Look at these extracts from a leaflet of Amoxicillin with Clavulanic Acid. Put the extracts into proper categories:

Instructions for Use	Undesirable Effects	Precautions
1, 3, 5, 8, 9, 12	4, 7, 10, 11,	2, 6,

1. I give you this drug in form of suspension, so take it orally
2. tell your doctor if you have or have ever had kidney or liver disease, allergies, asthma, hay fever, hives, or mononucleosis.
3. take with a meal or a snack
4. stop using if there are hives, skin rash or itching
5. take every 8 hours
6. inform your doctor if you are pregnant, plan to become pregnant, or are breast-feeding
7. yellowing of the skin and eyes may occur
8. do not take a double dose to make up for a missed one

9. take it around the same time every day
10. this drug may decrease the effectiveness of oral contraceptives so plan to use another form of birth control while taking amoxicillin and clavulanic acid
11. call your doctor immediately if you experience wheezing or breathing difficulty
12. shake the liquid well before each use to mix the medication evenly