

VOCABULARY PRACTICE – *KNOCKED-OUT TOOTH. PART TWO – ANSWERS*

Complete the text below with words from the table and then put the steps of replantation procedure in order

Background:

Extraoral dry time less than 60 min. The tooth has been kept in milk for less than 60 minutes.

1. Clean the root surface and apical foramen with a 1) **stream** of saline and soak the tooth in saline thereby removing 2) **contamination** and dead cells from the root surface.
6. Suture gingival lacerations if 3) **present**.
2. Administer 4) **local** anesthesia.
9. 5) **Administer** systemic antibiotics. Tetracycline is the first choice.
3. Irrigate the 6) **socket** with saline.
8. Apply a flexible splint for up to 2 weeks, keep 7) **away** from the gingiva.
4. Examine the alveolar socket. If there is a fracture of the socket wall, 8) **reposition** it with a suitable instrument.
5. Replant the tooth slowly with slight digital pressure. Do not use 9) **force**.
10. If the avulsed tooth has been in contact with 10) **soil**, and if tetanus coverage is uncertain, refer to physician for a tetanus 11) **booster**.
7. Verify normal position of the replanted tooth both, clinically and 12) **radiologically**.
11. Initiate root canal treatment 7-10 days after replantation and before splint removal.

Rearrange the words to create pieces of advice you would give your patient following avulsed tooth replantation

1. /to/take/up/weeks/soft/only/food/two/

Take only soft food up to two weeks.

2. (0.1 %)/twice/a chlorhexidine/one/a day/use/rinse/for/mouth/week

Rinse mouth twice a day with a chlorhexidine (0.1%) for one week.

3. 1 year/and then yearly thereafter/radiographic/6 months/schedule/after/clinical and/four weeks/control/3 months/

Schedule clinical and radiographic control four weeks, 3 months, six months and then yearly thereafter.

4. immediately/an antibiotic-corticosteroid/following/apply/paste/or shortly/replantation

Apply an antibiotic-corticosteroid paste immediately or shortly following replantation.

5. radiographic/I suggest/2 weeks/control/after

I suggest radiographic control after 2 weeks.

Find terms for the following definitions (all words are taken from this lesson) and complete the crossword puzzle

1. The forcible tearing away of a tooth due to trauma, accident or injury.
2. A dental appliance put on the teeth to protect them from moving out of place.
3. A tooth adapted for cutting or gnawing, located at the front of the mouth along the apex of the dental arch.
4. Adjective describing localization of something. Synonym of next to, nearby.
5. A solution that is isotonic with blood and is used in medicine and surgery.
6. An acute, often fatal disease characterized by spasmodic contraction of voluntary muscles, especially those of the neck and jaw, and caused by the toxin of the bacillus *Clostridium tetani*, which typically infects the body through a deep wound. Also called *lockjaw*.
7. The condition of being impure.
8. To reattach surgically to the original site.

1	A	V	U	L	S	I	O	N					
2	S	P	L	I	N	T							
3	I	N	C	I	S	O	R						
4	A	D	J	A	C	E	N	T					
5	S	A	L	I	N	E							
6	T	E	T	N	U	S							
7	C	O	N	T	A	M	I	N	A	T	I	O	N
8	R	E	P	L	A	N	T	A	T	I	O	N	